Supporting Information

Evolution of the Snow Area Index (SAI) of the subarctic snowpack in central Alaska over a whole season. Consequences for the air to snow transfer of pollutants.

A.-S. Taillandier[1,2], F. Domine[1,2*], W.R. Simpson[2,3], M. Sturm[4], T.A. Douglas[4], and K. Severin[5]

[1] CNRS, Laboratoire de Glaciologie et Géophysique de l'Environnement, BP 96, 38402 Saint-Martin d'Hères Cedex, France
[2] Geophysical Institute, University of Alaska Fairbanks, Alaska, USA
[3] Department of Chemistry, University of Alaska Fairbanks, USA
[4] U.S. Army Cold Regions Research and Engineering Laboratory, Fort Wainwright, Alaska, USA
[5] Department of Geology and Geophysics, University of Alaska Fairbanks, USA
* corresponding author: florent@lgge.obs.ujf-grenoble.fr

1 Figure



Figure S1. Evolution of crystal morphology of the snow layer fallen on 10 November. Scale bars: 1 mm (except for (c) where it is 200 µm). (a) Fresh dendritic crystal. (b) Decomposing dendritic crystal a few hours later. (c) One and a half day old faceted crystal, seen with an ESEM. (d) One week old striated crystal. (e), (f) and (g) Depth hoar crystals of increasing age: 3, 5 and 12 weeks, respectively.